

INTRAVENEOUS IMMUNE GLOBULINS

BRAND NAME	Gammagard S/D		IGIVnex & GAMUNEX®	Gammagard Liquid	Privigen®	Panzyga®
WEBLINK	http://www.baxalta.ca/products_en.html		www.grifols.com	http://www.baxalta.ca/products_en.html	www.cslbehring.ca	www.octapharma.ca
SIZES	5g		5g, 10g, 20g Only 20g for IGIVnex	2.5g, 5g, 10g, 20g, 30g	2.5g, 5g, 10g, 20g, 40g	10g, 20g, 30g
INDICATIONS	PID, CLL, ITP		PID, SID, ITP, CIDP, GBS	PID, SID, ITP, MMN	PID, SID, ITP, CIDP	PID, SID, ITP
CONCENTRATION	5% or 10% upon reconstitution		10%	10%	10%	10%
IgA CONTENT	≤ 2.2 µg/mL in 5% solution		Not more than 0.084mg/mL	≤ 0.14 mg/mL	No more than 25 mcg/mL	≤0.3 mg/mL
SUGAR CONTENT	20 mg/mL dextrose in 5%		Not indicated in product monograph	None	Not indicated in product monograph	Not indicated in product monograph
STORAGE TEMPERATURE	Not to exceed +25°C		+2°C to +8°C	+2 to +8°C	+2°C to +25°C	+2°C to +8 °C
INITIAL Recommended INFUSION RATE	<u>5% & 10% Solution</u> 0.5 mL/kg of body weight/hr		0.01 to 0.02 mL/kg/ min (1 to 2 mg/kg/min) for the first 30 mins	0.5 mL/kg of BW/hr for 30 mins BW=body weight	0.5 mg/kg/min (0.3 mL/kg/hr)	1 mg/kg/ min for the first 30 mins
MAXIMUM Recommended INFUSION RATE	<u>5% Solution</u> 4 mL/kg/hr	<u>10% Solution</u> 8 mL/kg/hr	0.14 mL/kg/min (14 mg/kg/min)	8 mL/kg of BW/hr <u>For MMN</u> 5.4 mL/kg of body weight/hr	12 mg/kg/min (7.2 mL/kg/hr)	8 mg/kg/min (in new patients) <u>For PID & SID</u> 12 or 14 mg/kg/min (in experienced patients)

2018-03-20

This Table is a summary and will not tell you everything about IG products. **DETAILED information can be found in the PRODUCT MONOGRAPH or PRODUCT INSERT.** *CONTENT is subject to change.*