SUBCUTANEOUS IMMUNE GLOBULINS TABLE

BRAND NAME	IGIVnex & GAMUNEX®1	Hizentra®2	Cuvitrus	cutaquig®4
WEBLINK	www.grifols.com	www.cslbehring.ca	www.shirecanada.com	www.octapharma.ca
INDICATIONS	Primary Immune Deficiency Secondary Immune Deficiency			
CONCENTRATION	10%	20%	20%	16.5%
SIZES	2.5g, 5g, 10g, 20g 20g only for IGIVnex	1g, 2g, 4g, 10g Prefilled syringes for 1g, 2g	1g, 2g, 4g, 8g	1g, 2g, 4g, 8g
IgA CONTENT	Not more than 0.084 mg/mL	No more than 50 mg/L	≤280 mcg/mL with an average concentration of 80 mcg/mL	Not indicated in PM
SUGAR CONTENT	Not indicated in Product Monograph	None	None	Not indicated in PM
STORAGE TEMPERATURE	+2°C to +8°C	+2°C to +25°C	+2°C to +8°C	+2°C to +8°C
RECOMMENDED INFUSION RATE	20 mL/hr per infusion site	20 mL/hr per site	10-20 mL/hr per site	25 mL/hr/site

Product Monograph (PM) reference: 1.IGIVnex PM July 2017 & GAMUNEX® PM Jan 2016, 2. Hizentra® PM Aug 2019, 3. Cuvitru PM July 2018, 4. Cutaquig PM Aug 2019